

Title	Abundance of σ -compact non-compactly generated groups witnessed by the Bohr topology of an abelian group (General and geometric topology today and their problems)
Author(s)	Dikranjan, Dikran; Shakhmatov, Dmitri
Citation	数理解析研究所講究録 (2008), 1578: 16-18
Issue Date	2008-02
URL	http://hdl.handle.net/2433/81381
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Abundance of σ -compact non-compactly generated groups witnessed by the Bohr topology of an abelian group *

Dikran Dikranjan ¹

Dipartimento di Matematica e Informatica
Università di Udine, Via delle Scienze 206, 33100 Udine, Italy

Dmitri Shakhmatov ²

Graduate School of Science and Engineering
Division of Mathematics, Physics and Earth Sciences
Ehime University, Matsuyama 790-8577, Japan

Abstract

Answering negatively a question of Fujita and Shakhmatov [2], Tkachenko and Torres Falcón [6] have constructed an example of a countable (and thus, σ -compact) totally bounded group that is not compactly generated. We observe that *any* countable non-finitely generated abelian group G equipped with its Bohr topology fails to be compactly generated, thereby obtaining an abundant supply of totally bounded groups providing a counter-example to the question of Fujita and Shakhmatov [2].

A group G is *finitely generated* if G is algebraically generated by its finite subset.

A group G is called *σ -compact* provided that G can be represented as a union of a countable family of its compact subsets, and G is called *compactly*

*This manuscript is in its final form and will not be submitted for publication elsewhere.

¹The first author was partially supported by MEC. MTM2006-02036 and FEDER FUNDS. *e-mail*: dikranja@dimi.uniud.it

²The second author was partially supported by the Grant-in-Aid for Scientific Research no. 19540092 by the Japan Society for the Promotion of Science (JSPS). *e-mail*: dmitri@dpc.ehime-u.ac.jp

generated if G is algebraically generated by its compact subset. One can easily see that a compactly generated group is σ -compact.

Fujita and Shakhmatov [2] proved that a σ -compact metric group is compactly generated. The same result also holds for a wider class of groups that contains both metric and locally compact groups, see [3].

Answering a question of Fujita and Shakhmatov [2], Tkachenko and Torres Falc3n [6] have constructed an example of a countable (thus, σ -compact) totally bounded group that is not compactly generated. (Recall that a group G is *totally bounded* if it is (topologically and algebraically) isomorphic to a subgroup of some compact group.)

The main purpose of this note is to observe that *any* countable non-finitely generated abelian group G equipped with its Bohr topology fails to be compactly generated, thereby obtaining an abundant supply of totally bounded groups providing a counter-example to the question of Fujita and Shakhmatov [2].

Let G be an abelian group. The *Bohr topology* of G is the weakest group topology on G making all characters $\chi : G \rightarrow \mathbb{T}$ continuous, where \mathbb{T} is the torus group. The Bohr topology of G is totally bounded, and the group G equipped with this topology is usually denoted by $G^\#$.

According to the celebrated result of Glicksberg [4], $G^\#$ has no infinite compact subsets. Thus, $G^\#$ is compactly generated if and only if $G^\#$ is finitely generated. This immediately yields the following

Theorem 1. *Let G be a countable abelian group that is not finitely generated. Then $G^\#$ is a σ -compact totally bounded group that is not compactly generated.*

It should be noted that there are only countably many finitely generated abelian groups, as every such group has the form

$$\mathbb{Z}^n \times \mathbb{Z}(m_1) \times \dots \times \mathbb{Z}(m_k), \quad (1)$$

where n, m_1, \dots, m_k are integer numbers and $\mathbb{Z}(k)$ denotes the cyclic group of order k . On the other hand, there are continuum many pairwise non-isomorphic countable abelian groups. (In fact, even the group \mathbb{Q} of rational numbers contains continuum many pairwise non-isomorphic subgroups.)

Corollary 2. *Let G be a countable abelian group that is not isomorphic to a group of the form (1). Then $G^\#$ is a σ -compact totally bounded group that is not compactly generated.*

Let us finish with some concrete examples.

Corollary 3. *$\mathbb{Q}^\#$ is a (σ -compact) divisible totally bounded group that is not compactly generated.*

Corollary 4. *If G is a countably infinite torsion group, then $G^\#$ is a $(\sigma$ -compact) totally bounded group that is not compactly generated.*

It follows from Corollary 4 that $(\mathbb{Q}/\mathbb{Z})^\#$ is a $(\sigma$ -compact) divisible totally bounded group that is not compactly generated.

Furthermore, from Corollary 4 one can easily obtain an infinite family of $(\sigma$ -compact) non-compactly-generated totally bounded torsion groups that are pairwise non-homeomorphic even as topological spaces. Indeed, if G and H are countably infinite torsion groups of distinct prime exponent, then $G^\#$ and $H^\#$ are not homeomorphic as topological spaces ([5, 1]).

References

- [1] D. Dikranjan and S. Watson, *A solution to van Douwen's problem on the Bohr topologies*, J. Pure Appl. Algebra **163** (2001), 147–158.
- [2] H. Fujita and D. Shakhmatov, *A characterization of compactly generated metric groups*, Proc. Amer. Math. Soc. **131** (2003), 953–961.
- [3] H. Fujita and D. Shakhmatov, *Topological groups with dense compactly generated subgroups*, Applied Gen. Topol. **3** (2002), 85–89.
- [4] I. Glicksberg, *Uniform boundedness for groups*, Canad. J. Math. **14** (1962), 269–276.
- [5] K. Kunen, *Bohr topology and partition theorems for vector spaces*, Topology Appl. **90** (1998), 97–107.
- [6] M. Tkachenko and Y. Torres Falc3n, *An example of a σ -compact monothetic group which is not compactly generated*, Bol. Soc. Mat. Mexicana (3) **10** (2004), no. 2 (2005), 203–208.